

1 February 2008

High-precision ϵ -expansions of massive four-loop vacuum bubbles

S. Laporta*

*Dipartimento di Fisica, Università di Bologna,
Via Irnerio 46, I-40126 Bologna, Italy*

Abstract

In this paper we calculate at high-precision the expansions in $\epsilon = (4 - D)/2$ of the master integrals of 4-loop vacuum bubble diagrams with equal masses, using a method based on the solution of systems of difference equations. We also show that the analytical expression of a related on-shell 3-loop self-mass master integral contains new transcendental constants made up of complete elliptic integrals of first and second kind.

PACS number(s):

12.20.Ds, 12.38.Bx

Keywords: Feynman diagram, master integral, difference equation.

*E-mail: laporta@bo.infn.it

$$\begin{aligned}
V_2 = & -1.6875\epsilon^{-4} - 7.8125\epsilon^{-3} - 21.209635416666666666666666667\epsilon^{-2} \\
& - 44.769552951388888888888888889\epsilon^{-1} - 97.0765348083978788434432266577 \\
& - 290.923474358239921908941634044\epsilon - 1719.80956172642828117680003592\epsilon^2 \\
& - 8934.7307852874907107364237238\epsilon^3 - 51529.1688840498204793229332431\epsilon^4 \\
& - 236942.747528861416676127442283\epsilon^5 - 1169551.39766202377654297049171\epsilon^6 \\
& - 4923196.24852687778836136788114\epsilon^7 + O(\epsilon^8) \quad (2)
\end{aligned}$$

$$\begin{aligned}
V_3 = & 1.625\epsilon^{-4} + 10.229166666666666666666666667\epsilon^{-3} \\
& + 33.4657920462820508294798181834\epsilon^{-2} + 83.6384721220072849057411859385\epsilon^{-1} \\
& + 142.482921255630165397059484622 + 73.8226346988098680135859408617\epsilon \\
& - 2215.06968731839973043724485006\epsilon^2 - 10483.810544518473050561125786\epsilon^3 \\
& - 75868.1741728976518970530475501\epsilon^4 - 266235.221240550768248796525565\epsilon^5 \\
& - 1562163.76406718387868426308061\epsilon^6 - 5089761.51470684069337522895124\epsilon^7 + O(\epsilon^8) \quad , \quad (3)
\end{aligned}$$

$$\begin{aligned}
V_4 = & 1.5\epsilon^{-4} + 9.5\epsilon^{-3} + 33.5\epsilon^{-2} + 59.8938292905212171438007855155\epsilon^{-1} \\
& - 6.77093494521947617804741496782 - 927.184486122109949342449134087\epsilon \\
& - 7607.58935727331640672412576731\epsilon^2 - 36043.9315389277686247446140344\epsilon^3 \\
& - 199692.43129042192811514839511\epsilon^4 - 784516.116603849522465732718633\epsilon^5 \\
& - 3938768.43031938169221908816973\epsilon^6 - 14490248.5601255703505458679593\epsilon^7 + O(\epsilon^8) \quad , \quad (4)
\end{aligned}$$

$$\begin{aligned}
V_5 = & -0.25\epsilon^{-4} - 2.333333333333333333333333333\epsilon^{-3} \\
& - 11.7695293784587392828307280615\epsilon^{-2} - 47.3155150442894117036246868531\epsilon^{-1} \\
& - 139.695412330272368658893294367 - 952.24968928016088147255137961\epsilon \\
& - 1559.77276310374672922449535587\epsilon^2 - 17433.2756155912579714527203251\epsilon^3 \\
& - 18039.431294353693419505748424\epsilon^4 - 300299.661276324759969342746297\epsilon^5 \\
& - 223362.891943298104957627580543\epsilon^6 - 5002799.01434332803606709511202\epsilon^7 + O(\epsilon^8) \quad , \quad (5)
\end{aligned}$$

$$\begin{aligned}
V_6 = & 2.375\epsilon^{-4} + 15.979166666666666666666666667\epsilon^{-3} \\
& + 55.370266375339080107172526045\epsilon^{-2} + 180.944159673951583547109549717\epsilon^{-1} \\
& + 496.624640289958679240601686338 + 2103.52942297240529902308231413\epsilon \\
& + 4385.91019392343391919235019438\epsilon^2 + 29880.7052368668975066622604109\epsilon^3 \\
& + 45750.0902911026047573732270476\epsilon^4 + 472601.136971716877928457965404\epsilon^5 \\
& + 573888.829426582538928155340449\epsilon^6 + 7686577.54749127119979436449377\epsilon^7 + O(\epsilon^8) \quad , \quad (6)
\end{aligned}$$

$$\begin{aligned}
V_7 = & -0.5\epsilon^{-4} - 4.416666666666666666666666667\epsilon^{-3} \\
& - 18.4682782839316233281953131334\epsilon^{-2} - 68.0334660202028428854058007578\epsilon^{-1} \\
& - 295.653297261799619866589394558 - 1111.65699468499496145662198062\epsilon \\
& - 4624.23997399795582698262827805\epsilon^2 - 18166.6470161776485664443287094\epsilon^3 \\
& - 72747.4433773622788330319854055\epsilon^4 - 294544.59963387669806853126121\epsilon^5 \\
& - 1152092.06523577988360701658787\epsilon^6 - 4748621.41279753209699735445982\epsilon^7 + O(\epsilon^8) \quad , \quad (7)
\end{aligned}$$

$$\begin{aligned}
V_8 = & -0.75\epsilon^{-4} - 6.25\epsilon^{-3} - 23.7024174258974349922929697001\epsilon^{-2} \\
& - 83.9541325752545325454615616904\epsilon^{-1} - 438.152668276175094650239945304 \\
& - 1151.7188201038655941347197846\epsilon - 7624.47573539137683288770528724\epsilon^2 \\
& - 16496.7827202671608148012112968\epsilon^3 - 127976.07440527009995741832362\epsilon^4 \\
& - 245588.151837894378828676455499\epsilon^5 - 2103545.82723681108089748244607\epsilon^6 \\
& - 3759879.03947407762892613884153\epsilon^7 + O(\epsilon^8) , \quad (8)
\end{aligned}$$

$$\begin{aligned}
V_9 = & 1.80308535473939142809960724227\epsilon^{-2} - 9.0439355503127513874520962153\epsilon^{-1} \\
& + 42.0771149080147860090872289595 - 185.506435563765087317986949292\epsilon \\
& + 794.690451268783109449998200836\epsilon^2 - 3340.85301017420853074405241137\epsilon^3 \\
& + 13860.9560593967975329182060014\epsilon^4 - 56963.5538585614643909090940607\epsilon^5 \\
& + 232474.430355572470669990540268\epsilon^6 - 943890.666478469607638755776088\epsilon^7 + O(\epsilon^8) , \quad (9)
\end{aligned}$$

$$\begin{aligned}
V_{10} = & 5.18463877571684963165682743229\epsilon^{-1} - 38.9591108409575465929423742741 \\
& + 216.540829715907801334069893492\epsilon - 1056.20279561819887173122551098\epsilon^2 \\
& + 4804.63905121999951090192731115\epsilon^3 - 20965.3713155251719483741628451\epsilon^4 \\
& + 89093.2196878747322139561202477\epsilon^5 - 372004.961361167636317053811767\epsilon^6 \\
& + 1534700.00918085457184730424284\epsilon^7 + O(\epsilon^8) , \quad (10)
\end{aligned}$$

$$\begin{aligned}
V_{11} = & -1.36872356982626095206640482818\epsilon^{-2} - 10.1337693663646787142726963524\epsilon^{-1} \\
& + 25.6153496410310292617132790055 - 334.538889614391947948784653628\epsilon \\
& + 984.338776062634546439769565907\epsilon^2 - 7183.20521761939021795215293201\epsilon^3 \\
& + 21442.9895505509641128293112066\epsilon^4 - 132391.959284919922968209921626\epsilon^5 \\
& + 396184.588583865634141091817143\epsilon^6 - 2278903.80544414662394641296011\epsilon^7 + O(\epsilon^8) , \quad (11)
\end{aligned}$$

$$\begin{aligned}
V_{12} = & 1.34894802170970895986445430292 - 8.34992524160927013803891299755\epsilon \\
& + 40.4425656120211378353957075462\epsilon^2 - 177.294826919557277843981859823\epsilon^3 \\
& + 742.428926688827707784575315859\epsilon^4 - 3039.49399009410599098768805594\epsilon^5 \\
& + 12302.4392626409424372470869128\epsilon^6 - 49506.0256176923061393145333901\epsilon^7 + O(\epsilon^8) , \quad (12)
\end{aligned}$$

$$\begin{aligned}
V_{13} = & 0.997672576874263051049093586736 - 5.83175720330362704111639113258\epsilon \\
& + 27.5586906484187171474221359635\epsilon^2 - 119.248620281553405809848697481\epsilon^3 \\
& + 495.915629767162238727475039832\epsilon^4 - 2022.83520871375744081923368263\epsilon^5 \\
& + 8171.6838591733877817993555444\epsilon^6 - 32850.412079699027325130879249\epsilon^7 + O(\epsilon^8) . \quad (13)
\end{aligned}$$

The master integrals V_j are defined as

$$V_j = (\pi^{D/2}\Gamma(1+\epsilon))^{-4} \int d^D k_1 d^D k_2 d^D k_3 d^D k_4 \frac{P_j}{Q_j} ,$$

where $P_2 = (k_1 \cdot k_2)^2$, $P_6 = (k_1 + k_2)^2$, $P_{11} = k_1 \cdot k_2$ and all other $P_j = 1$; Q_j is the product of denominators of the corresponding j th diagram of Fig.1, each with unit mass. The expressions of the finite integrals V_{12} and V_{13} have been checked by comparing them with values obtained by performing Monte-Carlo integrations over Feynman parameters.

The rapid growth of the coefficients is due to the fact that the integrals, seen as functions of D , have poles near $D = 4$. For example, in the case of V_{13} the nearest poles are $D = 9/2, 5, 16/3$, and $11/2$; by factorizing out these poles, one obtains a series with decreasing coefficients:

$$V_{13} = \left[(1 + 4\epsilon)(1 + 2\epsilon) \left(1 + \frac{3}{2}\epsilon \right) \left(1 + \frac{4}{3}\epsilon \right) \right]^{-1} \left(.9976726 + 2.981017\epsilon + 2.981995\epsilon^2 \right. \\ \left. + 1.315019\epsilon^3 + .432643\epsilon^4 + .0999339\epsilon^5 + .0150665\epsilon^6 + .0086152\epsilon^7 + O(\epsilon^8) \right). \quad (14)$$

Now we sketch here the method used for obtaining Eqs.(1)-(13); for a complete description the reader is referred to Refs.[20, 21]. For each master integral V_j we have chosen a denominator and raised it to n , and we have built a system of difference equations between the functions $V_j(n)$. The difference equation for a given master integral contains in the r.h.s. only master integrals with less denominators, which are simpler. The equations of the system are solved one at once, beginning with that corresponding to the simplest master integral, V_1 , and ending with that corresponding to the most complex, V_{13} . The necessary boundary conditions at $n \rightarrow \infty$ are expressed by 3-loop integrals which are calculated by solving other systems of difference equations. The homogeneous and nonhomogeneous solutions of the difference equations are expanded in factorial series or alternatively transformed into integral representations, and evaluated numerically for suitable integer values of the exponents. By taking these results as input values and by using the equations of the system as recurrence relations, one recovers the values of the master integrals $V_j(1)$, Eqs.(1)-(13).

For example, the function $V_1(n)$ satisfies the fourth-order difference equation

$$p_1 V_1(n+1) + p_2 V_1(n) + p_3 V_1(n-1) + p_4 V_1(n-2) + p_5 V_1(n-3) = 6(D-2)^3 J^3(1) J(n-2), \quad (15)$$

$$\begin{aligned} p_1 &= -90n(n-1)(n-2), \quad p_2 = 9(n-1)(n-2)(16n-9D-6), \\ p_3 &= (n-2)(-20n^2 + (122-21D)n + 27D^2 - 66D - 42), \\ p_4 &= -32n^3 + (54+93D)n^2 + (-78D^2 - 153D + 14)n + 18D^3 + 87D^2 + 21D - 24, \\ p_5 &= -(n-D-1)(n-2D+1)(2n-3D). \end{aligned}$$

It contains in the r.h.s. the integral obtained from V_1 by contracting a line, which factorizes into a product of 4 one-loop tadpoles. The solution of Eq.(15) compatible with the large- n boundary condition $V_1(n) \propto n^{-D/2}$ can be written as $V_1(n) = C I^{HO}(n) + I^{NH}(n)$, where I^{HO} and I^{NH} are respectively the solution of the homogeneous equation with above large- n behaviour and a particular solution of the nonhomogeneous equation (15). The constant C turns out to be the value of the 3-loop vacuum bubble obtained from V_1 by deleting a line.

The amount of calculations needed to work out and solve all the systems of difference equations is rather high, so that the calculations have been performed by means of an automatic tool, the program **SYS** described in Ref.[20].

Eqs.(1)-(13) have a precision sufficient to substitute analytical expressions for most of practical uses. Anyway, it is interesting to work out the analytical expressions of some of these results. In the case of V_4 , a straightforward application of the methods used in [17] for calculating the 3-loop g -2

master integrals gives

$$V_4 = \frac{3}{2}\epsilon^{-4} + \frac{19}{2}\epsilon^{-3} + \frac{67}{2}\epsilon^{-2} + \left(-3\zeta(3) + \frac{127}{2}\right)\epsilon^{-1} - \frac{\pi^4}{20} + 97\zeta(3) - \frac{237}{2} + \epsilon\left(189\zeta(5) - \frac{559}{60}\pi^4 - 32\pi^2\ln^2 2 + 768\text{Li}_4\left(\frac{1}{2}\right) + 32\ln^4 2 + 1261\zeta(3) - \frac{3969}{2}\right) + O(\epsilon^2) , \quad (16)$$

which agrees perfectly with Eq.(4). As usual, the expression turns out to contain integer values of Riemann ζ -function $\zeta(p) = \sum_{n=1}^{\infty} 1/n^p$, and values of the polylogarithms $\text{Li}_p(x) = \sum_{n=1}^{\infty} x^n/n^p$. The same method may be also used for calculating V_8 , but not for calculating the remaining master integrals. The reason is the appearance of double elliptic integrals which cannot be expressed in terms of polylogarithms.

The appearance of elliptic integrals in the calculation of multiloop diagrams with many massive lines is a well-known phenomenon. In fact, the principal difficulty of the analytical calculation of 3-loop g -2 has been the individuation of the approaches which allowed us to avoid elliptic integrations. Unfortunately in the case of the 4-loop vacuum master integrals the problem cannot be circumvented and we are forced to perform the elliptic integrations.

Let us consider the simplest vacuum diagram, V_1 . By cutting one line of this bubble one obtains the 3-loop self-mass diagram shown in Fig.2. If we take all masses equal to one, and the external line on the mass-shell, this diagram has two master integrals, whose S_1 is

$$S_1 = (\pi^{D/2}\Gamma(1+\epsilon))^{-3} \int d^D k_1 d^D k_2 d^D k_3 \frac{1}{Q} ,$$

and Q is the product of the four denominators of the diagram.

The analytical expressions of V_1 and S_1 are strictly related. In fact, introducing hyperspherical variables[25, 26] and performing the angular integrations one finds that the finite parts of V_1 and S_1 can be expressed as a sum of many double elliptic integrals with the same three square roots in the denominator. The simplest of these integrals is that with unity as numerator

$$A = \int_0^\infty \frac{dl}{R(l, -1, -1)} \int_0^\infty \frac{dm}{R(m, l, -1)R(m, -1, -1)} = 2.641379476074689\dots, \quad (17)$$

$$R(x, y, z) = \sqrt{x^2 + y^2 + z^2 - 2xy - 2xz - 2yz} . \quad (18)$$

We have been not able to calculate this integral in analytical form. Therefore, we have resorted to ‘experimental mathematics’: we have evaluated it at very high precision and we have tried to fit the numerical value with various kinds of analytical expressions. Luckily¹, we have found that

$$A = K(w_-)K(w_+) , \quad w_\pm = \frac{z_\pm}{z_\pm - 1} , \quad z_\pm = -(2 - \sqrt{3})^4(4 \pm \sqrt{15})^2 , \quad (19)$$

¹ We have been inspired by the analytical expressions of a double elliptic integral[27] related to the Appell F_2 function and by the analytical result of the 2-loop 3-lines “sunrise” self-mass diagram in two dimensions

$$S(p^2 = -1, m_1^2 = 1, m_2^2 = 1 - 2^{-1/2}, m_3^2 = 1 + 2^{-1/2}, D = 2) = \pi K(1/2)/2 = \sqrt{\pi}\Gamma^2(1/4)/8 .$$

where K is the first of the two standard elliptic integrals

$$K(m) = \int_0^1 \frac{dt}{\sqrt{1-t^2}\sqrt{1-mt^2}}, \quad E(m) = \int_0^1 \frac{dt \sqrt{1-mt^2}}{\sqrt{1-t^2}}. \quad (20)$$

We have verified Eq.(19) with a precision of more than 30000 digits². Once the analytical expression of the basic integral A has been identified beyond any reasonable doubt, we have considered the full integral representations of the finite parts of V_1 and S_1 ; they contain integrals which differ from Eq.(17) by additional polynomials and logarithmic functions in the numerators. A high-precision numerical expansion in ϵ of S_1 was calculated in Ref.[28] by means of the program **SYS**. We have been able to fit the numerical result with the analytical expression

$$S_1(\text{Fig.2}) = 2\epsilon^{-3} + \frac{22}{3}\epsilon^{-2} + \frac{577}{36}\epsilon^{-1} + \frac{4\pi}{\sqrt{15}} \left(\frac{35}{8}\pi + \frac{131}{12}K(w_-)K(w_+) \right. \\ \left. - \frac{7}{2}(E(1-w_-)E(1-w_+) + 5E(w_-)E(w_+)) \right) + \frac{6191}{216} + O(\epsilon) \quad (21)$$

by using the integer-relation search algorithm PSLQ[29]. We have verified Eq.(21) with 1200 digits of precision. Eq.(21) is important because it shows for the first time that there exist single-scale integrals which contain non-polylogarithmic transcendental constants. The same constants also appear in the analytical expression of V_1 , together with higher-transcendentality constants so far unidentified.

This result also sheds light on the possible analytical structure of the 4-loop QED contribution to the electron $g-2$. In fact one of the 4-loop $g-2$ master integrals is a 4-loop 5-lines self-mass integral analogous to S_1 , expressible in terms of irreducible triple elliptic integrals analogous to Eq.(17).

References

- [1] K. G. Chetyrkin and F. V. Tkachov, *Nucl. Phys.* **B192** (1981) 159.
- [2] F. V. Tkachov, *Phys. Lett.* **B100** (1981) 65.
- [3] S. G. Gorishny, *Nucl. Phys.* **B319** (1989) 633.
- [4] K. G. Chetyrkin, *Teor. Mat. Fiz.* **75** (1988) 26; *ibid.* **76** (1988) 207.
- [5] V. A. Smirnov, *Commun. Math. Phys.* **134** (1990) 109.
- [6] V. A. Smirnov, *Renormalization and asymptotic operations* (Birkhäuser,Basel,1991).
- [7] M. Beneke and V. A. Smirnov, *Nucl. Phys.* **B522** (1998) 321.
- [8] V. A. Smirnov, *Mod.Phys.Lett.* **A10** (1995) 1485.
- [9] R. Harlander and M. Steinhauser, *Prog.Part.Nucl.Phys.* **43** (1999) 167.
- [10] M. Steinhauser, *Phys.Rept.* **364** (2002) 247.

² By making the replacement $R(l, -1, -1) \rightarrow R(l, -1, \lambda)$, working out the differential equation in λ satisfied by $A(\lambda)$, and evaluating $A(1)$ from the series expansion of the solution, with the initial condition $A(0) = \pi^2/4$.

- [11] K. G. Chetyrkin and M. Steinhauser, *Nucl. Phys.* **B573** (2000) 617.
- [12] J. Fleischer, M. Yu. Kalmykov and A. V. Kotikov, *Phys. Lett.* **B462** (1999) 169.
- [13] J. Fleischer and M. Yu. Kalmykov, *Phys. Lett.* **B470** (1999) 168.
- [14] M. Yu. Kalmykov and O. Veretin, *Phys. Lett.* **B483** (2000) 315.
- [15] A. I. Davydychev and M. Yu. Kalmykov, *Nucl. Phys.* **B605** (2001) 266.
- [16] D. Broadhurst, *Eur. Phys. J.* **C8** (1999) 311.
- [17] S. Laporta and E. Remiddi, *Phys. Lett.* **B379** (1996) 283.
- [18] S. Laporta and E. Remiddi, *Acta Phys. Polon.* **B28** (1997) 959.
- [19] K. Melnikov and T. V. Ritbergen, *Nucl. Phys.* **B591** (2000) 515.
- [20] S. Laporta, *Int. J. Mod. Phys.* **A15** (2000) 5087.
- [21] S. Laporta, *Phys. Lett.* **B504** (2001) 188.
- [22] S. Laporta, *Phys. Lett.* **B523** (2001) 95.
- [23] A. V. Kotikov, *Phys. Lett.* **B254** (1991) 158.
- [24] E. Remiddi, *Nuovo Cimento* **A110** (1997) 1435.
- [25] M. J. Levine and R. Roskies, *Phys. Rev.* **D9** (1974) 421.
- [26] M. J. Levine, E. Remiddi and R. Roskies, in: *Quantum Electrodynamics*, edited by T. Kinoshita, Advanced series on Directions in High Energy Physics, Vol. 7 (World Scientific, Singapore, 1990) p. 162.
- [27] A. P. Prudnikov, Yu. A. Brychov, O. I. Marichev, in *Integrals and Series*, vol.1: *Elementary Functions*, Gordon and Breach, New York, 1992, p. 575, eq. 3.1.5.16.
- [28] See Ref.[20], Eq.205, integral $I(2e)$.
- [29] H. R. P. Ferguson and D. H Bailey, RNR Technical Report RNR-91-032.

Figure Captions

Figure 1: Four-loop vacuum bubble diagrams.

Figure 2: Three-loop self-mass diagram.

(S. Laporta, High-precision ϵ -expansions of massive four-loop vacuum bubbles)




Figure 1:

(S. Laporta, High-precision ϵ -expansions of massive four-loop vacuum bubbles)




Figure 2:

(S. Laporta, High-precision ϵ -expansions of massive four-loop vacuum bubbles)